arXiv:gr-qc/0010039v2 24 Oct 2000

General exact solution of the Einstein–Dirac equations with the cosmological constant in homogeneous space

V. A. Zhelnorovich*
Institute of Mechanics, Moscow state university, Michurinsky pr., 1, Moscow 119899, Russia.

The general exact solution of the Einstein–Dirac equations with cosmological constant in the homogeneous Riemannian space of the Bianchi 1 type is obtained.

04.40.Nr, 04.20.Jb

I. INTRODUCTION

While integrating the Einstein–Dirac equations, one has to face two difficulties.

The first one is a purely technical difficulty related to the Einstein–Dirac equations being a complicated system of nonlinear partial differential equations of the second order with 24 unknown functions. Some particular exact solutions[1−6] of the Einstein-Dirac equations in homogeneous spaces used to be obtained before only for the diagonal metrics of Riemannian space.

The second difficulty is of a fundamental nature and related to the spinor field in Riemannian space being definable only in some nonholonomic orthonormal bases (tetrads) to be set. In other words, a gauge of the tetrads is necessary. Lots of such gauges are known, and different authors proposed different gauges. Taken together these gauges either are noninvariant with respect to transformation of variables in the observer's coordinate system or are written in the form of differential equations, which results in a complication of the initial system of equations.

Physically, all gauges are equivalent, since the Einstein-Dirac equations are tetrad–choice invariant. Mathematically, the use of a bad gauge (i.e. additional equations closing the Einstein-Dirac ones) may seriously complicate the equations, but the use of a good gauge may significantly simplify them.

To the difficulty in a reasonable gauge of tetrads detect is much related that in previous papers[1−6] solutions of the Einstein–Dirac equations have been obtained only for diagonal metrics (for such metrics the vectors of the basis of a holonomic coordinate system are orthogonal, and hence it is possible to make a natural choice of the tetrads related with an orthogonal holonomic basis of Riemannian space).

Here we use the tetrad gauge[7] that is algebraic and, at the same time, is formulated in an invariant way. Using this gauge allows one to reduce the number of unknown functions in the Einstein–Dirac equations by six units conserving the invariance of equations with respect to transformation of variables of the observer's coordinate system.

With the tetrad gauge being used here, all equations are formulated as the first-order equations only for two invariants of the spinor field and Ricci rotation coefficients of the proper bases defined by the spinor field. The Dirac equations after the tetrad gauge transform into equations for Ricci rotation coefficients and for invariants of the spinor field, with the Ricci rotation coefficients entering in these equations linearly and derivative-free. Therefore in a homogeneous Riemannian space the Dirac equations prove to close the Einstein equations for the Ricci rotation coefficients without using additional equations. In this case one may first integrate the first order equations for the Ricci rotation coefficients and the spinor field invariants, and then integrate the first order equations for scale factors.

With these two circumstances — a reduction of the number of unknown functions by six units and a possibility of integrating the second order equations in two steps — considerable

---

*Electronic mail: zhelnor@inmech.msu.su

simplification of the Einstein–Dirac equations is related, which makes it possible to obtain new exact solutions of these equations.

II. SPINOR FIELDS IN THE FOUR-DIMENSIONAL RIEMANNIAN SPACE

Let V is the four-dimensional Riemannian space with the metric signature $(+,+,+,-)$, referred to a system of coordinate with the variables x^i and the holonomic vector basis $\mathcal{J}_i$, $i=1,\,2,\,3,\,4$. The metric tensor of the space V is determined in basis $\mathcal{J}_i$ by covariant components g_{ij}; the connection is determined by the Christoffel symbols Γ^s_{ij}. Let us introduce in the space V a smooth field of the orthonormal bases (tetrads) $\mathbf{e}_a(x^i)$ $(a=1,\,2,\,3,\,4)$, by the relations

$$\mathbf{e}_a = h^i{}_a \mathcal{J}_i, \qquad \mathcal{J}_i = h_i{}^a \mathbf{e}_a, \tag{1.1}$$

where $h_i{}^a$, $h^i{}_a$ are the scale factors. We designate the indices of tensor components in the basis $\mathcal{J}_i$ by Latin letters $i,\ j,\ k,\ ...$; the indices of tensor components in the orthonormal basis $\mathbf{e}_a$ will be designated by the first letters of the Latin alphabet $a,\ b,\ c,\ d,\ e,\ f$.

The differential of the orthonormal basis vectors $\mathbf{e}_a(x^i)$ is defined by the Ricci rotation coefficients

$$d\mathbf{e}_a = \Delta_{i,a}{}^b \mathbf{e}_b dx^i, \tag{1.2}$$

which are expressed via scale factors

$$\Delta_{i,ac} = \frac{1}{2}\big[h^j{}_c\big(\partial_i h_{ja} - \partial_j h_{ia}\big) - h^j{}_a\big(\partial_i h_{jc} - \partial_j h_{ic}\big) + h_i{}^b h^j{}_a h^s{}_c\big(\partial_j h_{sb} - \partial_s h_{jb}\big)\big]. \tag{1.3}$$

Here $\partial_i = \partial/\partial x^i$ is the symbol of a partial derivative with respect to the variable x^i.

Let us determine in the Riemannian space V a spinor field of the first rank $\psi(x^i)$, given by contravariant components $\psi^A(x^i)$ $(A=1,\,2,\,3,\,4)$ in the orthonormal bases $\mathbf{e}_a(x^i)$. The spinor indeces of the components of any spin-tensor are lowered or raised by the formulas

$$\psi^A = e^{AB}\psi_B, \qquad \psi_A = e_{AB}\psi^B, \tag{1.4}$$

where $E = \|e_{AB}\|$, $E^{-1} = \|e^{AB}\|$ are the covariant and contravariant components of the metric spinor, defined by the equations

$$\gamma_a^T = -E\gamma_a E^{-1}, \qquad E^T = -E. \tag{1.5}$$

Here T is the transposition symbol; γ_a are the four-dimensional Dirac matrices that on definition satisfy the equation

$$\gamma_a\gamma_b + \gamma_b\gamma_a = 2g_{ab}I, \tag{1.6}$$

where I is the unit four-dimensional matrix, $\|g_{ab}\| = \mathrm{diag}(1,1,1,-1)$ the covariant components of metric tensor in an orthonormal basis $\mathbf{e}_a$.

It should be noted, that in the equations (1.4) contracting is always done with respect to the second index of antisymmetric components of the metric spinor.

Let us define also the conjugate spinor field with the covariant components $\psi^+ = \|\psi^+_A\|$ by the relation $\psi^+ = \dot{\psi}^T\beta$, where the point above a letter denotes the complex conjugation; the invariant spinor of second rank β is defined by the equations

$$\dot{\gamma}_a^T = -\beta\gamma_a\beta^{-1}, \qquad \dot{\beta}^T = \beta. \tag{1.7}$$

The four-component spinor field ψ in general case has two real invariants ρ, η, which can be defined by the equation

$$\rho\exp i\eta = \psi^+\psi + i\psi^+\gamma^5\psi, \tag{1.8}$$

where $\gamma^5 = \frac{1}{24}\varepsilon^{abcd}\gamma_a\gamma_b\gamma_c\gamma_d$, ε^{abcd} are the components of the four-dimensional Levi–Chivita tensor $\varepsilon^{1234} = -1$. In the Riemannian space V via the spinor field ψ and conjugate spinor field ψ^+ it is possible to define the proper orthonormal basis $\breve{\mathbf{e}}_a$ of the spinor field

$$\breve{\mathbf{e}}_1 = \pi^i\mathcal{J}_i, \qquad \breve{\mathbf{e}}_2 = \xi^i\mathcal{J}_i, \qquad \breve{\mathbf{e}}_3 = \sigma^i\mathcal{J}_i, \qquad \breve{\mathbf{e}}_4 = u^i\mathcal{J}_i. \tag{1.9}$$

The vector components π^i, ξ^i, σ^i, u^i in the equations (1.9) are defined by the relations[8,9]

$$\begin{aligned} \rho\pi^i &= \mathrm{Im}(\psi^T E\gamma^i\psi), \\ \rho\xi^i &= \mathrm{Re}(\psi^T E\gamma^i\psi), \\ \rho\sigma^i &= \psi^+\gamma^i\gamma^5\psi, \\ \rho u^i &= i\psi^+\gamma^i\psi, \end{aligned} \tag{1.10}$$

where the invariant ρ is determined by the relation (1.8), the spin-tensors $\gamma^i = h^i{}_a\gamma^a$ satisfy the equation

$$\gamma^i\gamma^j + \gamma^j\gamma^i = 2g^{ij}I. \tag{1.11}$$

It is obvious, that the scale factors $\breve{h}^i{}_a$, corresponding to the proper basis $\breve{\mathbf{e}}_a$, are determined by matrix

$$\breve{h}^i{}_a = \begin{Vmatrix} \pi^1 & \xi^1 & \sigma^1 & u^1 \\ \pi^2 & \xi^2 & \sigma^2 & u^2 \\ \pi^3 & \xi^3 & \sigma^3 & u^3 \\ \pi^4 & \xi^4 & \sigma^4 & u^4 \end{Vmatrix}. \tag{1.12}$$

If the spin-tensors γ_a, E, β are determined by matrices

$$\begin{gathered} \gamma_1 = \begin{Vmatrix} 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ -i & 0 & 0 & 0 \end{Vmatrix}, \qquad \gamma_2 = \begin{Vmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{Vmatrix}, \qquad \gamma_3 = \begin{Vmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \\ -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{Vmatrix}, \\ \gamma_4 = \begin{Vmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{Vmatrix}, \qquad E = \begin{Vmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{Vmatrix}, \qquad \beta = \begin{Vmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{Vmatrix}, \end{gathered} \tag{1.13}$$

then the components of spinor ψ calculated in the proper basis $\breve{\mathbf{e}}_a$, are determined by the invariants ρ, η as follows[10]

$$\begin{aligned} \breve{\psi}^1 &= 0, \qquad \breve{\psi}^2 = i\sqrt{\tfrac{1}{2}\rho}\,\exp\left(\tfrac{i}{2}\eta\right), \\ \breve{\psi}^3 &= 0, \qquad \breve{\psi}^4 = i\sqrt{\tfrac{1}{2}\rho}\,\exp\left(-\tfrac{i}{2}\eta\right). \end{aligned} \tag{1.14}$$

As is well known, the covariant derivative of spinor fields ψ, ψ^+ in the Riemannian space are defined by the relations[11]

$$\nabla_s\psi = \partial_s\psi - \frac{1}{4}\Delta_{s,ij}\gamma^i\gamma^j\psi, \qquad \nabla_s\psi^+ = \partial_s\psi^+ + \frac{1}{4}\psi^+\Delta_{s,ij}\gamma^i\gamma^j. \tag{1.15}$$

For the covariant derivative of spinor fields ψ, ψ^+ the equations are valid

$$\begin{aligned} \nabla_s\psi &= \left(-\frac{1}{4}\breve{\Delta}_{s,ij}\gamma^i\gamma^j + \frac{1}{2}I\partial_s\ln\rho - \frac{1}{2}\gamma^5\partial_s\eta\right)\psi, \\ \nabla_s\psi^+ &= \psi^+\left(\frac{1}{4}\breve{\Delta}_{s,ij}\gamma^i\gamma^j + \frac{1}{2}I\partial_s\ln\rho - \frac{1}{2}\gamma^5\partial_s\eta\right). \end{aligned} \tag{1.16}$$

Here the invariants of spinor field ρ, η are defined by the equation (1.8), the Ricci rotation coefficients $\breve{\Delta}_{i,jk} = \breve{h}_j{}^b\breve{h}_k{}^c\breve{\Delta}_{i,bc}$ correspond to the proper bases $\breve{\mathbf{e}}_a$ of a spinor field and are calculated by the formula

$$\breve{\Delta}_{s,ij} = \frac{1}{2}\big(\pi_i\nabla_s\pi_j - \pi_j\nabla_s\pi_i + \xi_i\nabla_s\xi_j - \xi_j\nabla_s\xi_i + \sigma_i\nabla_s\sigma_j - \sigma_j\nabla_s\sigma_i - u_i\nabla_s u_j + u_j\nabla_s u_i\big). \tag{1.17}$$

The relations (1.16) in the four-dimensional pseudoeuclidean space have been obtained in[12,13]. The relations (1.16) in a Riemannian space can be obtained from the corresponding relations in pseudoeuclidean space by replacement particular derivative to covariant ones. The relations (1.16) are satisfied identically in view of definitions of the quantities ρ, η, $\breve{\Delta}_{s,ij}$.

III. EINSTEIN–DIRAC EQUATIONS WITH THE COSMOLOGICAL CONSTANT

Let us consider the set of equations

$$\begin{aligned} \gamma^a\nabla_a\psi + m\psi &= 0, \\ R_{ab} - \frac{1}{2}Rg_{ab} + \lambda g_{ab} &= \kappa T_{ab}. \end{aligned} \tag{2.1}$$

Here ψ is a four-component spinor field in the four-dimensional Riemannian space of events V, given in the orthonormal basis $\mathbf{e}_a$; m, λ, κ are constants; $R = g^{ab}R_{ab}$ is the scalar curvature of space V, R^{ab} the components of the Ricci tensor calculated in basis $\mathbf{e}_a$ and $g^{ab} = \mathrm{diag}(1, 1, 1, -1)$.

The components T_{ab} of the energy–momentum tensor of the spinor field in basis $\mathbf{e}_a$ are defined by the relation

$$T_{ab} = \frac{1}{4}\left(\psi^+\gamma_a\nabla_b\psi - \nabla_b\psi^+\cdot\gamma_a\psi + \psi^+\gamma_b\nabla_a\psi - \nabla_a\psi^+\cdot\gamma_b\psi\right). \tag{2.2}$$

The equations (2.1) are invariant under the arbitrary pseudoorthogonal transformations of tetrads $\mathbf{e}_a$, therefore for the closure of equations (2.1) it is necessary to add the additional equations, defining tetrads $\mathbf{e}_a$. Such additional equations usually are called as the gauge conditions. Further on we accept the gauge conditions $\mathbf{e}_a = \breve{\mathbf{e}}_a$, i. e. we accept, that the arbitrary tetrad $\mathbf{e}_a$ in the equations (2.1) coincides with the proper tetrad of the spinor field ψ. In this case the scale factors $h^i{}_a$ in the equations (2.1) coincide with coefficients $\breve{h}^i{}_a$, determined by matrix (1.12).

The Dirac equations in this case are written as the equations on the Ricci rotation coefficients $\breve{\Delta}_{a,bc}$ and the invariants of the spinor field[7]

$$\begin{aligned} &\breve{D}_a \ln\rho + \breve{\Delta}_{b,a}{}^b = 2m\breve{\sigma}_a\sin\eta, \\ &\breve{D}^a\eta + \frac{1}{2}\varepsilon^{abcd}\breve{\Delta}_{b,cd} = 2m\breve{\sigma}^a\cos\eta. \end{aligned} \tag{2.3}$$

Here $\breve{D}_a = \breve{h}^i{}_a\partial_i = \{\pi^i\partial_i, \xi^i\partial_i, \sigma^i\partial_i, u^i\partial_i\}$; $\breve{\sigma}_a = \breve{\sigma}^a = (0, 0, 1, 0)$ are the components of the vector $\breve{\mathbf{e}}_3$ in the proper basis.

Coefficients $\breve{\Delta}_{a,bc}$ in the equations (2.3) are connected with the scale factors $\breve{h}^i{}_a$ by relation

$$\breve{\Delta}_{a,bc} = \frac{1}{2}\big[\breve{h}^j{}_a(\breve{D}_b\breve{h}_{jc} - \breve{D}_c\breve{h}_{jb}) + \breve{h}^j{}_c(\breve{D}_a\breve{h}_{jb} + \breve{D}_b\breve{h}_{ja}) - \breve{h}^j{}_b(\breve{D}_a\breve{h}_{jc} + \breve{D}_c\breve{h}_{ja})\big]. \tag{2.4}$$

The equations (2.3) are the identical record of the Dirac equations in the proper basis $\breve{\mathbf{e}}_a$. These equations can be received also from the Dirac equations in (2.1), if to replace in them derivatives $\nabla_i\psi$ via the formula (1.16) and to produce algebraic transformations.

The replacement in equations (2.3) coefficients $\breve{\Delta}_{a,bc}$ via $\breve{h}^i{}_a$ yields the following system of the invariant tensor equations[7]

$$\begin{aligned} &\nabla_i\rho\pi^i = 0, \qquad \nabla_i\rho\xi^i = 0, \qquad \nabla_i\rho\sigma^i = 2m\rho\sin\eta, \qquad \nabla_i\rho u^i = 0, \\ &\nabla^i\eta - \frac{1}{2}\varepsilon^{ijms}\left(\pi_j\nabla_m\pi_s + \xi_j\nabla_m\xi_s + \sigma_j\nabla_m\sigma_s - u_j\nabla_m u_s\right) = 2m\sigma^i\cos\eta. \end{aligned} \tag{2.5}$$

The Einstein equations in the proper basis $\breve{\mathbf{e}}_a$ it is convenient to write as

$$\breve{R}_{ab} = \kappa\breve{T}_{ab} + \left(\frac{1}{2}\kappa m\rho\cos\eta + \lambda\right)g_{ab}. \tag{2.6}$$

For transformation of the Einstein equations to the form (2.6) it is necessary to take into account, that in view of the equations (2.1) the follow equation is valid

$$R = -\kappa T_a{}^a + 4\lambda = \kappa m\rho\cos\eta + 4\lambda. \tag{2.7}$$

The expression of tetrad components of the energy–momentum tensor in gauge $\mathbf{e}_a = \breve{\mathbf{e}}_a$ is obtained in[7]:

$$\breve{T}_{ab} = \frac{1}{4}\rho\left[-\breve{\sigma}_b\breve{D}_a\eta - \breve{\sigma}_a\breve{D}_b\eta + \frac{1}{2}\breve{\sigma}_e\left(\breve{\Delta}_{a,cd}\varepsilon_b{}^{cde} + \breve{\Delta}_{b,cd}\varepsilon_a{}^{cde}\right)\right]. \tag{2.8}$$

The tensor components $\breve{R}_{ab}$ we can express via the Ricci rotation coefficients

$$\breve{R}_{ab} = \frac{1}{\sqrt{-g}}\partial_j\left[\sqrt{-g}\left(\breve{h}^j{}_c\breve{\Delta}_{b,a}{}^c - \breve{h}^j{}_b\breve{\Delta}_{c,a}{}^c\right)\right] - \breve{\Delta}_{f,b}{}^c\breve{\Delta}_{c,a}{}^f + \breve{\Delta}_{c,a}{}^c\breve{\Delta}_{f,b}{}^f. \tag{2.9}$$

Here $g = \det\|g_{ij}\|$.

Let us give also the expression of the Ricci rotation coefficients $\breve{\Delta}_{a,bc}$ immediately via the vector components π_i, ξ_i, σ_i, u_i, which is obtained from the formula (2.4), if replace the components $\breve{h}^i{}_a$ by the formula (1.12)

$$\begin{aligned}
&\breve{\Delta}_{1,12} = \xi^i\breve{D}_1\pi_i - \pi^i\breve{D}_2\pi_i, \qquad \breve{\Delta}_{2,12} = -\pi^i\breve{D}_2\xi_i + \xi^i\breve{D}_1\xi_i,\\
&\breve{\Delta}_{3,12} = \tfrac{1}{2}\big(\xi^i\breve{D}_3\pi_i - \pi^i\breve{D}_3\xi_i\big) + \tfrac{1}{2}\big(-\sigma^i\breve{D}_2\pi_i - \pi^i\breve{D}_2\sigma_i + \sigma^i\breve{D}_1\xi_i + \xi^i\breve{D}_1\sigma_i\big),\\
&\breve{\Delta}_{4,12} = \tfrac{1}{2}\big(\xi^i\breve{D}_4\pi_i - \pi^i\breve{D}_4\xi_i\big) + \tfrac{1}{2}\big(-u^i\breve{D}_2\pi_i - \pi^i\breve{D}_2u_i + u^i\breve{D}_1\xi_i + \xi^i\breve{D}_1u_i\big),\\
&\breve{\Delta}_{1,23} = \tfrac{1}{2}\big(\sigma^i\breve{D}_1\xi_i - \xi^i\breve{D}_1\sigma_i\big) + \tfrac{1}{2}\big(\pi^i\breve{D}_2\sigma_i + \sigma^i\breve{D}_2\pi_i - \pi^i\breve{D}_3\xi_i - \xi^i\breve{D}_3\pi_i\big),\\
&\breve{\Delta}_{2,23} = \sigma^i\breve{D}_2\xi_i - \xi^i\breve{D}_3\xi_i, \qquad \breve{\Delta}_{3,23} = -\xi^i\breve{D}_3\sigma_i + \sigma^i\breve{D}_2\sigma_i,\\
&\breve{\Delta}_{4,23} = \tfrac{1}{2}\big(\sigma^i\breve{D}_4\xi_i - \xi^i\breve{D}_4\sigma_i\big) + \tfrac{1}{2}\big(\sigma^i\breve{D}_2u_i + u^i\breve{D}_2\sigma_i - \xi^i\breve{D}_3u_i - u^i\breve{D}_3\xi_i\big),\\
&\breve{\Delta}_{1,31} = -\sigma^i\breve{D}_1\pi_i + \pi^i\breve{D}_3\pi_i, \qquad \breve{\Delta}_{3,31} = \pi^i\breve{D}_3\sigma_i - \sigma^i\breve{D}_1\sigma_i,\\
&\breve{\Delta}_{2,31} = \tfrac{1}{2}\big(\pi^i\breve{D}_2\sigma_i - \sigma^i\breve{D}_2\pi_i\big) + \tfrac{1}{2}\big(\pi^i\breve{D}_3\xi_i + \xi^i\breve{D}_3\pi_i - \xi^i\breve{D}_1\sigma_i - \sigma^i\breve{D}_1\xi_i\big),\\
&\breve{\Delta}_{4,31} = \tfrac{1}{2}\big(\pi^i\breve{D}_4\sigma_i - \sigma^i\breve{D}_4\pi_i\big) + \tfrac{1}{2}\big(\pi^i\breve{D}_3u_i + u^i\breve{D}_3\pi_i - \sigma^i\breve{D}_1u_i - u^i\breve{D}_1\sigma_i\big),\\
&\breve{\Delta}_{1,14} = u^i\breve{D}_1\pi_i - \pi^i\breve{D}_4\pi_i, \qquad \breve{\Delta}_{4,14} = -\pi^i\breve{D}_4u_i + u^i\breve{D}_1u_i,\\
&\breve{\Delta}_{2,14} = \tfrac{1}{2}\big(u^i\breve{D}_2\pi_i - \pi^i\breve{D}_2u_i\big) + \tfrac{1}{2}\big(\xi^i\breve{D}_1u_i + u^i\breve{D}_1\xi_i - \pi^i\breve{D}_4\xi_i - \xi^i\breve{D}_4\pi_i\big),\\
&\breve{\Delta}_{3,14} = \tfrac{1}{2}\big(u^i\breve{D}_3\pi_i - \pi^i\breve{D}_3u_i\big) + \tfrac{1}{2}\big(\sigma^i\breve{D}_1u_i + u^i\breve{D}_1\sigma_i - \pi^i\breve{D}_4\sigma_i - \sigma^i\breve{D}_4\pi_i\big),\\
&\breve{\Delta}_{1,24} = \tfrac{1}{2}\big(u^i\breve{D}_1\xi_i - \xi^i\breve{D}_1u_i\big) + \tfrac{1}{2}\big(\pi^i\breve{D}_2u_i + u^i\breve{D}_2\pi_i - \pi^i\breve{D}_4\xi_i - \xi^i\breve{D}_4\pi_i\big),\\
&\breve{\Delta}_{2,24} = u^i\breve{D}_2\xi_i - \xi^i\breve{D}_4\xi_i, \qquad \breve{\Delta}_{4,24} = -\xi^i\breve{D}_4u_i + u^i\breve{D}_2u_i,\\
&\breve{\Delta}_{3,24} = \tfrac{1}{2}\big(u^i\breve{D}_3\xi_i - \xi^i\breve{D}_3u_i\big) + \tfrac{1}{2}\big(\sigma^i\breve{D}_2u_i + u^i\breve{D}_2\sigma_i - \xi^i\breve{D}_4\sigma_i - \sigma^i\breve{D}_4\xi_i\big),\\
&\breve{\Delta}_{1,34} = \tfrac{1}{2}\big(u^i\breve{D}_1\sigma_i - \sigma^i\breve{D}_1u_i\big) + \tfrac{1}{2}\big(\pi^i\breve{D}_3u_i + u^i\breve{D}_3\pi_i - \pi^i\breve{D}_4\sigma_i - \sigma^i\breve{D}_4\pi_i\big),\\
&\breve{\Delta}_{2,34} = \tfrac{1}{2}\big(u^i\breve{D}_2\sigma_i - \sigma^i\breve{D}_2u_i\big) + \tfrac{1}{2}\big(\xi^i\breve{D}_3u_i + u^i\breve{D}_3\xi_i - \xi^i\breve{D}_4\sigma_i - \sigma^i\breve{D}_4\xi_i\big),\\
&\breve{\Delta}_{3,34} = u^i\breve{D}_3\sigma_i - \sigma^i\breve{D}_4\sigma_i, \qquad \breve{\Delta}_{4,34} = -\sigma^i\breve{D}_4u_i + u^i\breve{D}_3u_i.
\end{aligned} \tag{2.10}$$

At an integration of the Einstein–Dirac equations it is useful to take into account also the matrix of the components of the energy–momentum tensor, which is obtained in correspondence with definition (2.8):

$$\breve{T}_{ab} = \frac{1}{4}\rho\left\|\begin{matrix}
2\breve{\Delta}_{1,24} & \breve{\Delta}_{2,24} - \breve{\Delta}_{1,14} & -\breve{D}_1\eta + \breve{\Delta}_{3,24} & \breve{\Delta}_{4,24} - \breve{\Delta}_{1,12}\\
\breve{\Delta}_{2,24} - \breve{\Delta}_{1,14} & -2\breve{\Delta}_{2,14} & -\breve{D}_2\eta - \breve{\Delta}_{3,14} & -\breve{\Delta}_{2,12} - \breve{\Delta}_{4,14}\\
-\breve{D}_1\eta + \breve{\Delta}_{3,24} & -\breve{D}_2\eta - \breve{\Delta}_{3,14} & -2\breve{D}_3\eta & -\breve{D}_4\eta - \breve{\Delta}_{3,12}\\
\breve{\Delta}_{4,24} - \breve{\Delta}_{1,12} & -\breve{\Delta}_{2,12} - \breve{\Delta}_{4,14} & -\breve{D}_4\eta - \breve{\Delta}_{3,12} & -2\breve{\Delta}_{4,12}
\end{matrix}\right\|. \tag{2.11}$$

The equations (2.3), (2.6), (2.9) — (2.11) in the given system of coordinate x^i represent a closed set of equations for determining the functions $\pi_i(x^j)$, $\xi_i(x^j)$, $\sigma_i(x^j)$, $u_i(x^j)$, $\rho(x^j)$, $\eta(x^j)$. The contravariant components of the metric tensor of the Riemannian space are expressed via $\pi_i(x^j)$, $\xi_i(x^j)$, $\sigma_i(x^j)$, $u_i(x^j)$ by the relation

$$g^{ij} = \breve{h}^i{}_a\breve{h}^j{}_b g^{ab} = \pi^i\pi^j + \xi^i\xi^j + \sigma^i\sigma^j - u^iu^j. \tag{2.12}$$

IV. GENERAL EXACT SOLUTION OF THE EINSTEIN–DIRAC EQUATIONS IN HOMOGENEOUS SPACE

Let us consider a four-dimensional Riemannian space referred to synchronous system of coordinates with variables x^i, in which on definition are valid the equations

$$g_{44} = g^{44} = -1, \qquad g_{4\alpha} = g^{4\alpha} = 0, \qquad \alpha = 1, 2, 3. \tag{3.1}$$

We shall seek the solutions of the equations (2.3), (2.5), (2.6), (2.9) — (2.11) in the synchronous system of coordinate in the supposition, that all unknown functions depend only from parameter $x^4 = t$. Thus, it is suposed, that the space V is the homogeneous Bianchi 1 type Riemannian space.

In this case the equations (2.5) are written as

$$\begin{gathered} \partial_4(\sqrt{-g}\,\rho\pi^4) = 0, \qquad \partial_4(\sqrt{-g}\,\rho\xi^4) = 0, \qquad \partial_4(\sqrt{-g}\,\rho u^4) = 0, \\ \partial_4(\sqrt{-g}\,\rho\sigma^4) = 2m\sqrt{-g}\,\rho\sin\eta, \qquad \partial_4\eta = -2m\sigma^4\cos\eta. \end{gathered} \tag{3.2}$$

The set of equations (3.2) and equation

$$g^{44} \equiv \pi^4\pi^4 + \xi^4\xi^4 + \sigma^4\sigma^4 - u^4u^4 = -1 \tag{3.3}$$

are the complete set for determining the quantities π^4, ξ^4, σ^4, u^4, η, $\rho\sqrt{-g}$. The general solution of the equations (3.2), (3.3) has the form

$$\begin{gathered} \frac{C_\rho}{\rho\sqrt{-g}} = \frac{\pi^4}{C_\pi} = \frac{\xi^4}{C_\xi} = \frac{u^4}{C_u} = \frac{1}{\sqrt{1 + C_\sigma^2\cos^2(2mt+\varphi)}}, \\ \sigma^4 = \frac{\varepsilon C_\sigma \sin(2mt+\varphi)}{\sqrt{1 + C_\sigma^2\cos^2(2mt+\varphi)}}, \qquad \exp i\eta = \varepsilon\frac{1 + iC_\sigma\cos(2mt+\varphi)}{\sqrt{1 + C_\sigma^2\cos^2(2mt+\varphi)}}, \end{gathered} \tag{3.4}$$

where φ, C_π, C_ξ, C_σ, $C_u \geq 1$, $C_\rho > 0$ are the integration constants; coefficient ε can take any of the two values $+1$ or -1. By virtue of the condition of synchronism (3.3) the constans C are connected by the relation

$$(C_\pi)^2 + (C_\xi)^2 + (C_\sigma)^2 - (C_u)^2 = -1. \tag{3.5}$$

Let us introduce the notation $h_a = \breve{h}^4{}_a = (\pi^4, \xi^4, \sigma^4, u^4)$. From the solution (3.4) we find

$$\{h_1, h_2, h_4\} = \frac{1}{\sqrt{1 + C_\sigma^2\cos^2(2mt+\varphi)}}\{C_\pi, C_\xi, C_u\}. \tag{3.6}$$

It is clear from here, that by virtue of solution (3.4) the directions of the three-dimensional vector with components h_1, h_2, h_4 does not depend on parameter t.

From definition (2.4) it follows, that the Ricci rotation coefficients $\breve{\Delta}_{a,bc}$ can be presented in the form

$$\breve{\Delta}_{a,bc} = \frac{1}{2}(h_b s_{ac} - h_c s_{ab} - h_a a_{bc}), \tag{3.7}$$

where by definition

$$\begin{gathered} s_{ab} = s_{ba} = \breve{h}^i{}_a\partial_4\breve{h}_{ib} + \breve{h}^i{}_b\partial_4\breve{h}_{ia}, \\ a_{ab} = -a_{ba} = \breve{h}^i{}_a\partial_4\breve{h}_{ib} - \breve{h}^i{}_b\partial_4\breve{h}_{ia}. \end{gathered} \tag{3.8}$$

The quantities h_a in the formula (3.7) are determined by solution (3.4) as the functions of parameter t, therefore the relation (3.8) expresses 24 dependent functions $\breve{\Delta}_{a,bc}$ only through 16 functions s_{ab}, a_{ab}.

By virtue of definition (3.8) the quantities s_{ab} in a synchronous system of the coordinates satisfy the identities

$$h^b s_{ab} = 0, \qquad s_a{}^a = 2\partial_4 \ln\sqrt{-g}. \tag{3.9}$$

From the first identity in (3.9) it follows, that the ten symmetrical components s_{ab} contains generally the six independent ones.

From the equations (2.3) it follows, that the antisymmetric quantities a_{ab} are determined by equality

$$a_{ab} = 4m\left[(\breve{\sigma}_a h_b - \breve{\sigma}_b h_a)\sin\eta - \varepsilon_{abcd}\breve{\sigma}^c h^d \cos\eta\right]. \tag{3.10}$$

Using the equations (3.9), (3.10) and definitions (2.9), (2.11) for T_{ab}, R_{ab} we can transform the Einstein equations (2.6) into the equivalent set of equations

$$\begin{gathered}\partial_4(\sqrt{-g}\,s_{ab}) - 2\sqrt{-g}\,\left(m\cos\eta + \tfrac{1}{8}\kappa\rho\right)\left(\varepsilon_{cefa}s_b{}^f + \varepsilon_{cefb}s_a{}^f\right)h^c\breve{\sigma}^e - \\ -2m\sqrt{-g}\,\left(h_a s_{bc} + h_b s_{ac}\right)\breve{\sigma}^c\sin\eta = \left(\kappa m\rho\sqrt{-g}\cos\eta + 2\lambda\sqrt{-g}\right)\left(g_{ab} + h_a h_b\right), \\ \left(s_a{}^a\right)^2 - s_{ab}s^{ab} = 8(\kappa m\rho\cos\eta + \lambda).\end{gathered} \tag{3.11}$$

The quantity $\rho\sqrt{-g}\cos\eta$ in the right part of the first equation (3.11) by virtue of the solution (3.4) is constant

$$\rho\sqrt{-g}\cos\eta = \varepsilon C_\rho. \tag{3.12}$$

The first equation in (3.11) is obtained by contracting of the Einstein equations in (2.6) with the tensor components $\delta^a_c + h_c h^a$ by the index a. The second equation in (3.11) is obtained by contracting of the Einstein equations in (2.6) with the components of tensor $g^{ab} + 2h^a h^b$ by the indices a, b.

The contracting of the equations (3.11) with g^{ab} by the indices a, b gives the equation

$$\partial_4\partial_4\sqrt{-g} = \frac{3}{2}\kappa m\varepsilon C_\rho + 3\lambda\sqrt{-g} \tag{3.13}$$

for definition of the quantity $\sqrt{-g}$.

If the cosmological constant λ is positive $\lambda > 0$, the solution of the equation (3.13) has the form

$$\sqrt{-g} = \frac{\varepsilon\kappa m}{2\lambda}C_\rho\left[-1 + f_1\sinh\left(\sqrt{3\lambda}\,t\right) + f_2\cosh\left(\sqrt{3\lambda}\,t\right)\right], \tag{3.14}$$

where f_1, f_2 are arbitrary constants.

If $\lambda < 0$, for $\sqrt{-g}$ is obtained

$$\sqrt{-g} = \frac{\varepsilon\kappa m}{2\lambda}C_\rho\left\{-1 + f\sin\left[\sqrt{-3\lambda}\,(t - t_\circ)\right]\right\}, \tag{3.15}$$

where f, t_0 are the integration constants.

The case $\lambda = 0$ has been considered in[14,15].

As a direction of the three-dimensional vector with components h_1, h_2, h_4 does not depend on parameter t, then by the constant Lorentz transformation of the basis vectors $\breve{\mathbf{e}}_1$, $\breve{\mathbf{e}}_2$, $\breve{\mathbf{e}}_4$ it is possible to transform the components h_1, h_2 to zero. The initial set of equations (2.3), (2.6), (2.9) – (2.11) is invariant under the arbitrary constant Lorentz transformation of the vectors $\breve{\mathbf{e}}_1$, $\breve{\mathbf{e}}_2$, $\breve{\mathbf{e}}_4$. Therefore without loss of generallity it is enough to consider a solution of the equations (2.3), (2.6), (2.9) - (2.11) only at $h_1 = h_2 = 0$. Under this condition the first equation in (3.11) can be written in the form

$$
\begin{aligned}
&\partial_4(s_{33}\sqrt{-g}\,) - 4m\sigma^4 \sin\eta\, s_{33}\sqrt{-g} = \big(\varepsilon\kappa m C_\rho + 2\lambda\sqrt{-g}\,\big)(u^4)^2,\\
&\partial_4(s_{23}\sqrt{-g}\,) - 2m\sigma^4 \sin\eta\, s_{23}\sqrt{-g} - \left(2m\cos\eta + \tfrac{1}{4}\kappa\rho\right)u^4 s_{13}\sqrt{-g} = 0,\\
&\partial_4(s_{13}\sqrt{-g}\,) - 2m\sigma^4 \sin\eta\, s_{13}\sqrt{-g} + \left(2m\cos\eta + \tfrac{1}{4}\kappa\rho\right)u^4 s_{23}\sqrt{-g} = 0,\\
&\partial_4(s_{11}\sqrt{-g}\,) + 2\left(2m\cos\eta + \tfrac{1}{4}\kappa\rho\right)u^4 s_{12}\sqrt{-g} = \varepsilon\kappa m C_\rho + 2\lambda\sqrt{-g},\\
&\partial_4(s_{22}\sqrt{-g}\,) - 2\left(2m\cos\eta + \tfrac{1}{4}\kappa\rho\right)u^4 s_{12}\sqrt{-g} = \varepsilon\kappa m C_\rho + 2\lambda\sqrt{-g},\\
&\partial_4(s_{12}\sqrt{-g}\,) - \left(2m\cos\eta + \tfrac{1}{4}\kappa\rho\right)u^4 (s_{11} - s_{22})\sqrt{-g} = 0,\\
&\partial_4(s_{14}\sqrt{-g}\,) + \left(2m\cos\eta + \tfrac{1}{4}\kappa\rho\right)u^4 s_{24}\sqrt{-g} - 2mu^4 \sin\eta\, s_{31}\sqrt{-g} = 0,\\
&\partial_4(s_{24}\sqrt{-g}\,) - \left(2m\cos\eta + \tfrac{1}{4}\kappa\rho\right)u^4 s_{14}\sqrt{-g} - 2mu^4 \sin\eta\, s_{23}\sqrt{-g} = 0,\\
&\partial_4(s_{34}\sqrt{-g}\,) - 2m\sigma^4 \sin\eta\, s_{34}\sqrt{-g} = u^4(\varepsilon\kappa m C_\rho \sigma^4 + 2m\sin\eta\, s_{33}\sqrt{-g}\,),\\
&\partial_4(s_{44}\sqrt{-g}\,) - 4mu^4 \sin\eta\, s_{34}\sqrt{-g} = \big(\varepsilon\kappa m C_\rho + 2\lambda\sqrt{-g}\,\big)(\sigma^4)^2.
\end{aligned}
\tag{3.16}
$$

The general solution for s_{ab} has the form:

$$
\begin{aligned}
s_{11} &= \rho u^4\left[-\frac{1}{3}N + \frac{2}{3C_\rho C_u}\partial_4\sqrt{-g} + \frac{1}{2}B\sin 2(\zeta+\beta)\right],\\
s_{22} &= \rho u^4\left[-\frac{1}{3}N + \frac{2}{3C_\rho C_u}\partial_4\sqrt{-g} - \frac{1}{2}B\sin 2(\zeta+\beta)\right],\\
s_{33} &= \frac{2}{3}\rho(u^4)^3\left(N + \frac{1}{C_\rho C_u}\partial_4\sqrt{-g}\right),\\
s_{44} &= \frac{2}{3}\rho u^4(\sigma^4)^2\left(N + \frac{1}{C_\rho C_u}\partial_4\sqrt{-g}\right),\\
s_{12} &= -\frac{1}{2}\rho u^4 B\cos 2(\zeta+\beta),\\
s_{34} &= \frac{2}{3}\rho\sigma^4(u^4)^2\left(N + \frac{1}{C_\rho C_u}\partial_4\sqrt{-g}\right),\\
s_{13} &= \frac{1}{2}\rho(u^4)^2 A\cos(\zeta+\alpha), \qquad s_{14} = \frac{1}{2}\rho u^4\sigma^4 A\cos(\zeta+\alpha),\\
s_{23} &= \frac{1}{2}\rho(u^4)^2 A\sin(\zeta+\alpha), \qquad s_{24} = \frac{1}{2}\rho u^4\sigma^4 A\sin(\zeta+\alpha),
\end{aligned}
\tag{3.17}
$$

where A, B, N, α, β are the arbitrary constants; the quantity $\sqrt{-g}$ in the equations (3.17) is defined by the equalities (3.14), (3.15); the quantity ζ is defined by the follow relation

$$
\zeta = \int\left(2m\cos\eta + \frac{1}{4}\kappa\rho\right)u^4\,dt = \varepsilon\,\mathrm{arctg}\left(\frac{\tan(2mt+\varphi)}{\sqrt{1+C_\sigma^2}}\right) + \frac{1}{4}\kappa\tau. \tag{3.18}
$$

The integral $\tau = \int \rho u^4 dt$ in the formula (3.18) depends from the value of cosmologic constant λ and will be calculated further.

Substituting in the second equation in (3.11) the components s_{ab} by the formulas (3.17), we obtain a connection between the integration constants A, B, N and f_1, f_2. In the case $\lambda > 0$ is obtained the following relation

$$
f_2^2 - f_1^2 = 1 - \frac{\lambda}{\kappa^2 m^2}C_u^2\left(\frac{1}{4}A^2 + \frac{1}{4}B^2 + \frac{1}{3}N^2\right) \le 1.
$$

It is obvious, that if $f_2^2 \ge f_1^2$, then the formula (3.14) for $\sqrt{-g}$ can be presented as

$$
\sqrt{-g} = \frac{\varepsilon\kappa m}{2\lambda}C_\rho\left\{-1 + f\cosh\!\big[\sqrt{3\lambda}\,(t-t_\circ)\big]\right\}, \qquad |f| \le 1, \tag{3.19a}
$$

where $t_\circ$, f are the arbitrary constants, $f^2 = f_2^2 - f_1^2$.

The singular points of the solution (3.19 a) are determined by the equation $f\cosh\bigl[\sqrt{3\lambda}\,(t-t_\circ)\bigr] = 1$. The solution has one singular point, if $f = 1$ and two singular points, if $0 < f < 1$. If $f \le 0$ (in this case $\varepsilon = -1$), the solution has no singular points.

If $f_2^2 < f_1^2$, for $\sqrt{-g}$ the equality is valid

$$\sqrt{-g} = \frac{\varepsilon\kappa m}{2\lambda} C_\rho \left\{-1 + f\sinh\bigl[\sqrt{3\lambda}\,(t - t_\circ)\bigr]\right\}, \tag{3.19b}$$

where $f^2 = f_1^2 - f_2^2$. In this case the solution has one singular point, defined by the equation $f\sinh\bigl[\sqrt{3\lambda}\,(t-t_\circ)\bigr] = 1$.

If $\lambda < 0$, the following formula for the integration constant f in (3.15) is obtained

$$f^2 = 1 - \frac{\lambda}{\kappa^2 m^2} C_u^2 \left(\frac{1}{4}A^2 + \frac{1}{4}B^2 + \frac{1}{3}N^2\right) \ge 1.$$

In this case there are indefinitely many singular points, are determined by the equation $\sqrt{-g} = 0$.

The condition $\sqrt{-g} > 0$ yields a restriction on the values of the integration constants and the domain of existence of the solution.

Using definition (1.14) and formulas (3.4), we can write out solution for the components of spinor field in the proper basis

$$\breve{\psi} = \pm \left\| \begin{array}{c} 0 \\ i\sqrt{\varepsilon C_\rho \dfrac{1 + iC_\sigma\cos(2mt+\varphi)}{2\sqrt{-g}}} \\ 0 \\ i\sqrt{\varepsilon C_\rho \dfrac{1 - iC_\sigma\cos(2mt+\varphi)}{2\sqrt{-g}}} \end{array} \right\|,$$

where $\sqrt{-g}$, depending on a sign λ, is defined by the formula (3.15) or formulas (3.19 a,b).

The equations (3.4), (3.10), (3.17) completely determine the Ricci rotation coefficients $\breve{\Delta}_{a,bc}$ by the formula (3.7) and represent the first integral of the equations (2.3), (2.5), (2.6), (2.9) — (2.11). For determining of the general solution of the equations (2.3), (2.5), (2.6), (2.9) — (2.11) it is now enough to integrate the equations (3.8), from which follows

$$\partial_4 \breve{h}_{ib} = \frac{1}{2}\bigl(s_{ab} + a_{ab}\bigr)\breve{h}_i{}^a. \tag{3.20}$$

In view of definition (1.12) the quantities $\breve{h}_i{}^a$ and found solutions (3.10), (3.17) for s_{ab}, a_{ab}, the equation (3.20) is possible to transform to the form

$$\begin{aligned} \frac{d}{d\tau}\left(u^4\sigma_j - \sigma^4 u_j\right) &= \frac{1}{4}A\bigl[\pi_j\cos(\zeta+\alpha) + \xi_j\sin(\zeta+\alpha)\bigr], \\ &+ \left(u^4\sigma_j - \sigma^4 u_j\right)\left(\frac{1}{3\sqrt{-g}}\frac{d}{d\tau}\sqrt{-g} + \frac{1}{3}N\right), \\ \frac{d}{d\tau}\bigl(\xi_j + i\pi_j\bigr) &= \bigl(\xi_j + i\pi_j\bigr)\left(\frac{1}{3\sqrt{-g}}\frac{d}{d\tau}\sqrt{-g} - \frac{1}{6}N - i\frac{2m}{\rho}\cos\eta\right), \\ &- \frac{i}{4}\bigl(\xi_j - i\pi_j\bigr)B\exp\bigl[-2i(\zeta+\beta)\bigr] + \frac{i}{4}A\bigl(u^4\sigma_j - \sigma^4 u_j\bigr)\exp[-i(\zeta+\alpha)]. \end{aligned} \tag{3.21}$$

The differential $d\tau$ in (3.21) is defined by equality $d\tau = \rho u^4 dt$. The set of equations (3.21) should be supplemented by the condition of synchronism

$$g_{4\alpha} \equiv \sigma_4\sigma_\alpha - u_4 u_\alpha = 0, \qquad \alpha = 1,2,3. \tag{3.22}$$

For the parameter τ we have

$$\tau = \int \rho u^4 dt = C_\rho C_u \int \frac{dt}{\sqrt{-g}}.$$

If the cosmological constant is positive $\lambda > 0$ and $\kappa m \neq 0$, a calculation by means of the equations (3.19 a,b) gives

$$\begin{aligned} \tau &= \frac{2\lambda C_u}{\varepsilon\kappa m} \int \frac{dt}{-1 + f \sinh\bigl[\sqrt{3\lambda}\,(t - t_\circ)\bigr]} \\ &= \frac{2\sqrt{\lambda}\,C_u}{\sqrt{3}\,\varepsilon\kappa m} \frac{1}{\sqrt{1+f^2}} \ln \left| \frac{f - \sqrt{f^2+1} + \tanh\frac{1}{2}\sqrt{3\lambda}\,(t - t_0)}{f + \sqrt{f^2+1} + \tanh\frac{1}{2}\sqrt{3\lambda}\,(t - t_0)} \right| + \tau_0 \end{aligned} \tag{3.23a}$$

and

$$\begin{aligned} \tau &= \frac{2\lambda C_u}{\varepsilon\kappa m} \int \frac{dt}{-1 + f \cosh\bigl[\sqrt{3\lambda}\,(t - t_\circ)\bigr]} \\ &= \frac{2\sqrt{\lambda}\,C_u}{\sqrt{3}\,\varepsilon\kappa m} \frac{1}{\sqrt{1-f^2}} \ln \left| \frac{f - 1 + \sqrt{1-f^2}\tanh\frac{1}{2}\sqrt{3\lambda}\,(t - t_0)}{f - 1 - \sqrt{1-f^2}\tanh\frac{1}{2}\sqrt{3\lambda}\,(t - t_0)} \right| + \tau_0, \end{aligned} \tag{3.23b}$$

where τ_0 is the arbitrary constant.

At $\lambda < 0$ $(f^2 > 1)$ by means of (3.15) we obtain

$$\begin{aligned} \tau &= \frac{2\lambda C_u}{\varepsilon\kappa m} \int \frac{dt}{-1 + f \sin\bigl[\sqrt{-3\lambda}\,(t - t_\circ)\bigr]} \\ &= \frac{2\sqrt{-\lambda}\,C_u}{\sqrt{3}\,\varepsilon\kappa m} \frac{1}{\sqrt{f^2-1}} \ln \left| \frac{\tan\frac{1}{2}\sqrt{-3\lambda}\,(t - t_0) - f - \sqrt{f^2-1}}{\tan\frac{1}{2}\sqrt{-3\lambda}\,(t - t_0) - f + \sqrt{f^2-1}} \right| + \tau_0. \end{aligned} \tag{3.24}$$

Let us change now the unknowns functions $(\pi_\lambda, \xi_\lambda, \sigma_\lambda, u_\lambda) \to (\pi^0_\lambda, \xi^0_\lambda, \theta_\lambda)$ in the equations (3.21)

$$\begin{aligned} \xi_\lambda + i\pi_\lambda &= \bigl(\xi^0_\lambda + i\pi^0_\lambda\bigr)\,(\sqrt{-g}\,)^{1/3} \exp\Bigl(-\tfrac{1}{6}N\tau - i\zeta\Bigr), \\ \sigma_\lambda &= \theta_\lambda(\sqrt{-g}\,)^{1/3} u^4 \exp\Bigl(-\tfrac{1}{6}N\tau\Bigr), \\ u_\lambda &= \theta_\lambda(\sqrt{-g}\,)^{1/3} \sigma^4 \exp\Bigl(-\tfrac{1}{6}N\tau\Bigr). \end{aligned} \tag{3.25}$$

Here ζ is defined by the relation (3.18), quantities σ^4, u^4 are defined by the solution (3.4), $\sqrt{-g}$ is determined by equalities (3.15) or (3.19 a, b). In outcome the condition of synchronism (3.22) is satisfied identically, and the set of equations (3.21) goes over into the set of linear equations with the constant coefficients:

$$\begin{aligned} \frac{d}{d\tau}\bigl(\xi^0_\lambda + i\pi^0_\lambda\bigr) &= \frac{i}{4}\kappa\bigl(\xi^0_\lambda + i\pi^0_\lambda\bigr) - \frac{i}{4}B\bigl(\xi^0_\lambda - i\pi^0_\lambda\bigr)e^{-2i\beta} + \frac{i}{4}A\theta_\lambda e^{-i\alpha}, \\ \frac{d}{d\tau}\theta_\lambda &= \frac{1}{4}A\bigl(\pi^0_\lambda \cos\alpha + \xi^0_\lambda \sin\alpha\bigr) + \frac{1}{2}N\theta_\lambda. \end{aligned} \tag{3.26}$$

At $j = 4$ the equations (3.21) are satisfied identically by virtue of conditions $h_1 = h_2 = 0$.

Using definition (2.12), we obtain the expression for the spatial components of metric tensor $g_{\alpha\beta}$ via the vector components π^0_α, ξ^0_α, σ^0_α:

$$g_{\alpha\beta} = \bigl(\sqrt{-g}\,\bigr)^{\frac{2}{3}} e^{-\frac{1}{3}N\tau}\bigl(\pi^0_\alpha\pi^0_\beta + \xi^0_\alpha\xi^0_\beta + \theta_\alpha\theta_\beta\bigr). \tag{3.27}$$

To the equations (3.26) for each value of an index $\lambda = 1, 2, 3$ corresponds the characteristic equation for an eigenvalue q

$$\det\left\{ \frac{1}{4} \left\| \begin{matrix} B\sin 2\beta & -B\cos 2\beta + \kappa & A\cos\alpha \\ -B\cos 2\beta - \kappa & -B\sin 2\beta & A\sin\alpha \\ A\cos\alpha & A\sin\alpha & 2N \end{matrix} \right\| - qI \right\} = 0,$$

which is possible to write as

$$2(N - 2q)(16q^2 + \kappa^2 - A^2 - B^2) + A^2[2N - B\sin 2(\alpha - \beta)] = 0. \tag{3.28}$$

The solution of the equation (3.28) in general case is determined by the Cardano formulas. A simple solution of this equation is obtained in the case $A = 0$ and $2N = B\sin 2(\alpha-\beta)$. A solution of equations (3.26) in the case $A = 0$ has been considered in[14,15].

Let us consider the case, when is valid the equation $2N = B\sin 2(\alpha-\beta)$, $A \neq 0$. In this case for the eigenvalues we have $q_1 = \frac{1}{2}N$, $q_2 = \frac{1}{4}\bigl(A^2+B^2-\kappa^2\bigr)^{1/2}$ and $q_3 = -\frac{1}{4}\bigl(A^2+B^2-\kappa^2\bigr)^{1/2}$. If $0 < A^2+B^2 < \kappa^2$, it is possible to write a solution of the equations (3.26) as follows

$$\begin{aligned}\xi^0_\lambda + i\pi^0_\lambda &= e^{-i\alpha}\{-AQ_\lambda e^{\frac{1}{2}N\tau} + [-(\kappa + Be^{2i(\alpha-\beta)})F_\lambda + 4i\Lambda G_\lambda]\cos\Lambda\tau \\ &+ [-(\kappa + Be^{2i(\alpha-\beta)})G_\lambda - 4i\Lambda F_\lambda]\sin\Lambda\tau\}, \\ \theta_\lambda &= [\kappa - B\cos 2(\alpha-\beta)]Q_\lambda e^{\frac{1}{2}N\tau} + A(F_\lambda\cos\Lambda\tau + G_\lambda\sin\Lambda\tau).\end{aligned} \tag{3.29}$$

Here $\Lambda = \frac{1}{4}\sqrt{\kappa^2-A^2-B^2}$; F_λ, G_λ, Q_λ are the integration constants connected by the relation

$$\varepsilon^{\alpha\beta\lambda}F_\alpha G_\beta Q_\lambda = (\kappa^2-A^2-B^2)^{-3/2}, \tag{3.30}$$

where $\varepsilon^{\alpha\beta\lambda}$ is the Levi-Chivita symbol. A relation (3.30) expresses the equality between the value of $\sqrt{-g}$, calculated from the solution (3.29), by its values (3.15), (3.19 a,b). For spatial components of the metric tensor by means of the formulas (3.27) the following expression is obtained

$$\begin{aligned}g_{\alpha\beta} &= (\sqrt{-g}\,)^{2/3}e^{-\frac{1}{3}N\tau}\Bigl\{Q_\alpha Q_\beta\{A^2 + [\kappa - B\cos 2(\alpha-\beta)]^2\}e^{N\tau} \\ &+ (F_\alpha G_\beta + F_\beta G_\alpha)[M\sin 2\Lambda\tau - 4\Lambda B\sin 2(\alpha-\beta)\cos 2\Lambda\tau] \\ &+ F_\alpha F_\beta[16\Lambda^2 + M + M\cos 2\Lambda\tau + 4\Lambda B\sin 2(\alpha-\beta)\sin 2\Lambda\tau] \\ &+ G_\alpha G_\beta[16\Lambda^2 + M - M\cos 2\Lambda\tau - 4\Lambda B\sin 2(\alpha-\beta)\sin 2\Lambda\tau] \\ &+ (F_\alpha Q_\beta + F_\beta Q_\alpha)2\kappa A e^{\frac{1}{2}N\tau}\cos\Lambda\tau \\ &+ (G_\alpha Q_\beta + G_\beta Q_\alpha)2\kappa A e^{\frac{1}{2}N\tau}\sin\Lambda\tau\Bigr\},\end{aligned} \tag{3.31}$$

where the contracted notation is entered

$$M = A^2 + B^2 + \kappa B\cos 2(\alpha-\beta).$$

The quantity $\sqrt{-g}$ in the right part of equality (3.31) is defined by the equations (3.19 a,b) or (3.15). It is obvious, that by the constant transformation of the variable x^1, x^2, x^3 it is always possible to transform components F_α, G_α, Q_α to the form

$$\begin{gathered}F_\alpha = \bigl\{(\kappa^2-A^2-B^2)^{-1/2}, 0, 0\bigr\}, \qquad G_\alpha = \bigl\{0, (\kappa^2-A^2-B^2)^{-1/2}, 0\bigr\}, \\ Q_\alpha = \bigl\{0, 0, (\kappa^2-A^2-B^2)^{-1/2}\bigr\}.\end{gathered} \tag{3.32}$$

The formula (3.31) at $A^2+B^2 \neq 0$ determine the oscillatory regime of an motion to singular points of the solution.

If $A^2+B^2 > \kappa^2$, then the trigonometric functions in (3.29) are replaced by the hyperbolic ones.

To the diagonal metric g_{ij} there corresponds the case, when the integration constants A, B in the equations (3.26) are equal to zero. In this case we have

$$\begin{aligned}\xi^0_\lambda + i\pi^0_\lambda &= \kappa(-F_\lambda + iG_\lambda)\exp\tfrac{i}{4}\kappa\tau, \\ \theta_\lambda &= \kappa Q_\lambda\exp\tfrac{1}{2}N\tau.\end{aligned} \tag{3.33}$$

Taking into account equality (3.27), we obtain the solution for the components of the metric tensor

$$g_{\alpha\beta} = \kappa^2\bigl(\sqrt{-g}\bigr)^{2/3}\left[e^{-\frac{1}{3}N\tau}\bigl(F_\alpha F_\beta + G_\alpha G_\beta\bigr) + e^{\frac{2}{3}N\tau}Q_\alpha Q_\beta\right]. \tag{3.34}$$

Here the quantity $\sqrt{-g}$ is defined by the formulas (3.15) or (3.19 a,b), and $e^{N\tau}$ is defined by the relations

$$\begin{aligned} e^{N\tau} &= \left(\frac{f - 1 + \sqrt{1-f^2}\tanh \frac{1}{2}\sqrt{3\lambda}\,(t-t_0)}{f - 1 - \sqrt{1-f^2}\tanh \frac{1}{2}\sqrt{3\lambda}\,(t-t_0)} \right)^{2\varepsilon} && \text{for} \quad \lambda > 0, \quad f_1^2 < f_2^2, \\ e^{N\tau} &= \left(\frac{f - \sqrt{f^2+1} + \tanh \frac{1}{2}\sqrt{3\lambda}\,(t-t_0)}{f + \sqrt{f^2+1} + \tanh \frac{1}{2}\sqrt{3\lambda}\,(t-t_0)} \right)^{2\varepsilon} && \text{for} \quad \lambda > 0, \quad f_1^2 > f_2^2, \\ e^{N\tau} &= \left(\frac{\tan \frac{1}{2}\sqrt{-3\lambda}\,(t-t_0) - f - \sqrt{f^2-1}}{\tan \frac{1}{2}\sqrt{-3\lambda}\,(t-t_0) - f + \sqrt{f^2-1}} \right)^{2\varepsilon} && \text{for} \quad \lambda < 0. \end{aligned} \tag{3.35}$$

If F_λ, G_λ, Q_λ to define by the relations

$$F_\alpha = \left\{\kappa^{-1}, 0, 0, \right\}, \qquad G_\alpha = \left\{0, \kappa^{-1}, 0\right\}, \qquad Q_\alpha = \left\{0, 0, \kappa^{-1}\right\}, \tag{3.36}$$

then the metric (3.33) is diagonal.

If $A = B = N = 0$, then

$$g_{\alpha\beta} = \kappa^2 \left(\sqrt{-g}\right)^{2/3} \left(F_\alpha F_\beta + G_\alpha G_\beta + Q_\alpha Q_\beta\right), \tag{3.37}$$

where $\sqrt{-g}$ is determined by the formulas (3.19 a), (3.15) and $f^2 = 1$.

Some particular solutions of the Einstein–Dirac equations with the cosmological constant in the homogeneous Riemannian space of the Bianchi 1 type with the diagonal metric have been obtained in[5,6] (without cosmological constant see[1–4,14,15]).

Let us consider the transformation of coordinates $(x^\alpha, t) \to (x^\alpha, \tau)$ defined by the equation $d\tau = \rho u^4 dt$. The function $\tau(t)$ is determined by relations (3.23 a,b), (3.24). Quantities $\sqrt{-g}\, g^{4i}$ in the transformed system of coordinate with variables x^α, τ have the form

$$(\sqrt{-g}\, g^{4\alpha})' = 0, \qquad (\sqrt{-g}\, g^{44})' = \text{const}.$$

From here follows, that in the system of coordinate with the variables x^α, τ is satisfied the harmonicities condition

$$\left[\partial_j (\sqrt{-g}\, g^{ij})\right]' = \frac{d}{d\tau} (\sqrt{-g}\, g^{4i})' = 0.$$

Thus, parameter τ is the time in the harmonic system of coordinate.

---

[1] C. J. Isham and J. E. Nelson, Phys. Rev. D **10**, 3226 (1974).

[2] C. J. Radford and A. N. Klotz, J. Phys. A: Math., Gen. **16**, 317 (1983).

[3] Patrick Wils, J. Math. Phys. **32**, 231 (1991).

[4] Yu. P. Rybakov, B. Saha and G. N. Shikin, Izv. Vuzov, Fizika No **7**, 40 (1994).

[5] V. G. Bagrov, A. D. Istomin and V. V. Obuckov, Gravitation and Cosmology **2**, 117 (1996).

[6] G. Platania and R. Rosania, Europhysics Letters **37**, 585 (1997).

[7] V. A. Zhelnorovich, Dokl. Acad. Nauk SSSR **296**, 571 (1987).

[8] F. Gürsey, Nuovo Cimento **5**, 784 (1957).

[9] T. Takabayasi, Nucl. Phys. **6**, 477 (1958).

[10] V. A. Zhelnorovich, *Theory of spinors and its application in physics and mechanics* (Moscow, Nauka, 1982).

[11] V. Fock, Zs. für Physik **57**, 216 (1929).

[12] V. A. Zhelnorovich, Dokl. Acad. Nauk SSSR **311**, 590 (1990).

[13] V. A. Zhelnorovich, Dokl. Rus. Acad. Nauk **322**, 507 (1992).

[14] V. A. Zhelnorovich, Dokl. Rus. Acad. Nauk **346**, 21 (1996).

[15] V. A. Zhelnorovich, Gravitation and Cosmology **2**, 109 (1996).